

KPCOM Respiratory System Lecture 1


03/28/2025 0810-0900

Mechanical Aspects of Obstructive and Restrictive Diseases




**HN Mayrovitz PhD
mayrovit@nova.edu
drmayrovitz.com**

Normal Complete Flow-Volume Loop



- As lung expands from RV the inward lung recoil increases so P_{REC} increases
- Chest wall outward recoil force decreases as thoracic volume gets closer to 0 stress
- These two forces work against increasing IFR
- But, as lung expands, AWR falls and favors air inflow
- PIFR is achieved due to these oppositely directed factors

PIFR Determinants



Obstructive vs. Restrictive Lung Disease

Basic Features

**Obstructive = Abnormal Increase in ???
Airway Resistance (R)**

**Restrictive = Abnormal Decrease in ???
Respiratory Compliance (C)**
→ More difficult to expand
→ Greater recoil force

Could (and do) have combinations – Mixed Disease

Obstructive Lung Disease

Key Features of Main Obstructive Lung Diseases

Chronic Obstructive Pulmonary Disease

(COPD) → Emphysema



Chronic Bronchitis


- Bronchial Inflammation
- Mucus
- Cough



- Alveolar-capillary wall destruction
- Loss of alveoli
- Increased air spaces
- More compliant but less alveolar recoil
- Airway collapse due to traction loss
- More difficult to expel air
- Mucus secretion
- ASM enhanced contraction

Asthma

- 1. Airway smooth muscle (ASM) **contraction** in response to **neuro and inflammatory mediators**
- 2. Airway wall thickening by **hyperplasia** or **hypertrophy**, **edema** and cellular infiltration
- 3. Airway obstruction by **mucus**, secretions and cellular debris





Obstructive Overview: COPD vs. Asthma



- Both considered due to **chronic respiratory tract inflammation**
- Both associated with **increased airway resistance**
- Asthma → Variable airflow limitation that is **usually reversible**
- COPD → Persistent airflow limitation **usually irreversible**

Parameter	Asthma	COPD
Symptoms	<ul style="list-style-type: none"> • Wheeze • Cough • Short of breath Variable – not usually progressive	<ul style="list-style-type: none"> • Short of breath • Cough • Mucus Persistent-Progressive
Onset	Usually Young	Usually > 40 years
Course	Variable – not usually progressive	Progressive
Bronchodilator response	Usually Good	Usually Good
Steroids Response	Usually Good	Usually Poor
Main Features	<ul style="list-style-type: none"> • Bronchoconstriction • Mast cell activation • Hyperresponsive ASM • Edema • Mucus Plugging 	<ul style="list-style-type: none"> • Emphysema • Mucus exudate • Small airway fibrosis • Edema • Distal airway enlargement and destruction
Main Airways Generations	<ul style="list-style-type: none"> • Larger ≥ 2 mm • (0-7) conducting zone 	<ul style="list-style-type: none"> • Smaller < 2 mm • Mostly respiratory zone



Asthma: *Increased Airway Resistance*




Asthma: *Increased Airway Resistance*




Asthma: *Increased Airway Resistance*



$SG_{AW} = \text{Specific Conductance} = \Delta G_{AW} / \Delta V = \text{SLOPE}$




Obstructive Lung Disease: Flow-Volume Changes



As compared to Normal


- PEFR reduced
- TLC increased
- RV increased
- FRC increased




- Extent of changes depends on amount of R_{AW} increase and if coexisting Emphysema

Emphysema vs. Normal

Advanced Emphysema



Normal Lung



White specs are blood vessels: low in Emphysema

Restrictive Lung Disease

Restrictive Diseases: *Restricts Lung Expansion*

Factors → “PAINT”

SITE → CAUSES

Pleural → Scarring or Effusion or fibrosis etc

Alveolar → Edema or Hemorrhage

Interstitial → Interstitial Lung Disease or Fibrosis


Neuromuscular → ALS or Myopathy

Thoracic/Extra-thoracic → Obesity or Ascites


Example Conditions

- Interstitial Fibrosis
+ alveolar fibrous tissue
Lung becomes stiffer
(-) compliance
Inspiration more difficult
- Allergic Alveolitis
Alvoli Wall Thickens
(-) compliance
- Pleural Effusion
Intrapleural Fluid buildup:
(-) compliance
Pleural fibrosis & + rigidity:
(-) compliance


Obstructive vs. Restrictive Compliance Abnormalities




Differential Effects on Lung Dynamic Work




Differential Forced Expiratory Flow-Volume Patterns



Forced Expiratory *Volume-Time* Test



Forced Expiratory *Volume-Time* Test




Factors and Reference Ranges To be considered for "Normal"

- Gender → Male > Female
- Age → Younger > Older
- Height → Taller > Shorter

	Mild	Moderate	Severe	Very Severe
FEV_1 (% predicted)	≥ 80%	50% to <80%	30% to <50%	<30%
Stages of COPD (All have $(FEV_1/FVC) < 0.7$)				

Differential Forced Expiratory Volume-Time Test

Normal-Obstructive-Restrictive



<http://www.cdc.gov/niosh/topics/spirometry/RefCalculator.html>

Total Work (O_2 Cost) is Minimum at some TV x RR


WORK COMPONENTS

Elastic work to expand lung (W_E) $\sim 1/\text{Compliance}$

- Larger TV \rightarrow more W_E since C decreases with V
- Larger TV \rightarrow less relative ADS (ADS/TV is less/breath)
- For same alveolar ventilation W_E decreases with +RR

Nonelastic ($W_{NE} \sim \text{Resistance}$)


- Energy is LOST to overcome airway and tissue resistances
- Larger RR \rightarrow Larger $dV/dt \rightarrow$ Larger W_{NE}
- For same alveolar ventilation W_{NE} increases with +RR



Total Work minimum at some RR

Two Clinical Points:

- Obstructive Disease: "slow and deep"
- Restrictive Disease: "rapid and shallow"



Hyperinflation: Quick Summary

Static Hyperinflation (*At rest*)

Recoil pressure reduced → + End Expiratory Lung Volume (EELV)

Chronic Obstructive Pulmonary Disease (COPD) especially moderate to severe EMPHYSEMA → Loss of elastic tissue → - recoil pressure
Generally, increase in Total Lung Capacity (TLC)

Dynamic Hyperinflation (*Exercise*)

Start inhalation before full exhalation completed

Potentially all levels of COPD including ASTHMA
→ Temporary and Variable in extent
→ Can occur with no increase in TLC

Main Factors Determining Extent:

- 1) Degree of airflow limitation
- 2) time available for exhalation (ΔT)


Dynamic Hyperinflation – “Air Trapping”

Can occur with any increase in airway R
(Airflow Limitation)

- To prevent “air trapping” need to exhale the volume (ΔV) that was just inhaled (TV)
- Volume actually removed (δV) depends on the air flow Q and the time (ΔT) as $\delta V = Q \times \Delta T$


$\uparrow R$ & $\downarrow P_{\text{Recoil}}$ \Rightarrow $\downarrow Q$
So TIME needed for exhalation \uparrow

If next inspiration starts prior to expiration to true FRC then air trapping!




Hyperinflation

Normal



Diaphragm

Hyperinflated



Flattening of the diaphragm

Chronic Obstructive Pulmonary Disease (COPD)

Especially moderate to severe EMPHYSEMA

- **Loss of elastic tissue**
- **Reduced recoil pressure**
- **Increased end expiratory lung volume**
- **Increase Total Lung Capacity (TLC)**

Dynamic Hyperinflation – Effects Summary

- May occur at rest but often manifest if increased ventilation demand
- Increased respiratory rate (RR) further **shortens available exp time**
- Further air trapping results and TV begins to be limited by now **diminishing inspiratory capacity (IC) due to rising FRC**
- Inspiratory **muscle load increases** → greater recoil at elevated volume
- **Increased work of inspiration** and oxygen cost of breathing
- Increasing amounts of **dyspnea**

10 Short Interactive Review Questions

1. If lung compliance decreases, what is the effect on the work of inspiration?
2. If surfactant production is low or absent, what will be the effect on work of inspiration?
3. What is the name given to lung volume at the end of quiet expiration?
4. As you inspire does intrapleural pressure increase or decrease?
5. As you inspire does translung pressure increase or decrease?
6. Is total respiratory compliance greater or less than lung compliance?
7. What are the components of airway-alveolar time constant?
8. During a normal respiration cycle, when is the air flow zero?
9. A person with emphysema would have a high or low lung recoil pressure?
10. A person with interstitial fibrosis would have a high or low lung recoil pressure?

End Respiration Physiology Lecture 1