

Changes in Tissue Water and Indentation Resistance of Lymphedematous Limbs Accompanying Low Level Laser Therapy (LLLT)


**Harvey N. Mayrovitz, PhD
College of Medical Sciences
Nova Southeastern University**

**Suzanne Davey, OTR/L, CLT-LANA
Healing Hands of Lymphatics
Ft. Lauderdale, Florida**

Does LLLT acutely affect tissue hardness and/or water content?


Does LLLT acutely affect tissue hardness and/or water content?

NLN 2008 – San Diego



Does LLLT acutely affect tissue hardness and/or water content?

Patients studied had clinically identified lymphedema and fibrosis (37 Arms 36 Legs)



Does LLLT acutely affect tissue hardness and/or water content?

Patients studied had clinically identified lymphedema and fibrosis (37 Arms 36 Legs)



1 Pre-LLLT Tx

5 min
LLLT



2 Post – LLLT Tx

51.2 ± 1.5
min



3 Post – MLD Tx


M
L
D

Measurements

- Girth → Tape measure
- Tissue Water → Tissue Dielectric Constant (TDC)
- Tissue Indentation Resistance → Force for a fixed indentation

Legs: Lymphedema due to Varied Etiologies


Males: 18 Females: 18



Arms: All Breast Cancer Treatment-Related Lymphedema All Unilateral N=37



Arms: All Breast Cancer Treatment-Related Lymphedema All Unilateral N=37



Girth

**Measured
with fixed
tension**




Tissue Dielectric Constant (TDC)



TDC ~ % H₂O
100% ≈ 78

Tissue Resistance (Force-Indentation)



LLLT 1 min @ each site



Results


Pre-Treatment Comparisons

Affected vs. Control

Arms → 37


Legs → 36

Limb Girth




Affected > Contralateral

Tissue Water



Affected > Contralateral

Tissue Force



Affected > Contralateral


Results

LLLT Treatment Effects


Pre - Tx → Post - Tx

Pre - Tx → Post - MLD


Tissue Force - Arms



Tissue Force - Legs



Tissue Water



Results

Active LLLT Treatment


Vs.

Sham Treatment


N= 17 Legs

Active vs. Sham

Force @ 4 mm or TDC



Active vs. Sham



Summary

- Rapid acute reduction in local tissue water and hardness with 5 minute treatment
- Occurs with active LLLT & with SHAM but greater reductions with ACTIVE
- Tissue water & hardness trend back toward pre-tx levels by the end of an MLD session
- Specific role & mechanism of low level laser light in observed changes is not yet known



Thanks for your Attention